INDEX TO VOLUME XVII

New species in blackface type

AAMODT, OLAF S., A study of growth habit and rust reaction in crosses between Marquis, Kota, and Kanred wheats, 573–609

Abacá, bunchy top and heart rot of, 255–257

Abavit "B," dust, for control of barley stripe, 176–178

Abstracts, of papers presented at the eighteenth annual meeting of the American Phytopathological Society, Philadelphia, Pa., December, 1926, 39–59; of papers read at eleventh annual meeting of Pacific Division of American Phytopathological Society, University of Nevada, Reno, Nevada, June, 1927, 745–748

Acacia, dealbata, extraction of tannin from, 142; decurrens, extraction of tannin from, 142; farnesiana, extraction of tannin from, 138; mollissima, extraction of tannin from, 142

Acanthorhynchus vaccinii, associated with storage rot of cranberries, 652–653

Acid agglutination, correlation in strains of Bacterium phaseoli sojense, 49

Ada Red, and other apple varieties, susceptibility to cedar rust, with special reference to twig infections, 541–543

Agallia sticticollis, transmission of curly top of sugar beet by, 407–408

Age, effect of, on tobacco mosaic virus, 515–517

Agglutination, acid, correlation in strains of Bacterium phaseoli sojense, 49; tests, Bacterium phaseoli, Bact. phaseoli sojense, and Bact. flaccumfaciens, 54

Agrilus anxius, bronze birch borer, 51

Alfalfa, wilt disease caused by Fusarium sp., 337–338; yellowing of, caused by leafhoppers, 39

ALLISON, R. V., see FARIS, J. A.

Allium cepa, new Alternaria disease of, 115–132

Alternaria allii, n. sp., cause of purple leaf spot of Allium cepa, 115–132; control of, 130; growth in culture, 119–124; life history of, 124–129

Alternaria, solani, from potato tubers, variation of strains of, 56; spp., connected with target canker of apples and pears, 736; spp., in corn nodes, 41

Amaranthus blitum (=A. lividus L.) infected by Blakeslea, 183

"Amarillamiento" (yellowing) root disease of coffee, 150

American Phytopathological Society, abstracts of papers presented at Philadelphia meeting, December 1926, 39–59; Pacific Division, abstracts of papers read at, 745–748; report of 18th annual meeting, 343–352; summer meeting of, 415, 836

American type culture collection, 672

Amphorophora rubi, aphid carrier of raspberry mosaic, 46

ANDERSON, H. W., A theory to account for the bactericidal action of sodium silicofluoride and lack of injury to host tissues, 50

Andes, central, plant disease notes from, 147–153

Anonymous, The American type culture collection, 672

Aphanomyces, cuteiches, 43; cuteiches, hosts of, 133–134; cuteiches, on Lathyrus latifolius, L. odoratus, L. sp., on Medicago sativa, Melilotus alba, Vicia angustifolia, V. dasycarpa, V. ervilia, V. fulgens, V. gigantea, V. monantha, V. pannonica, V. sativa, 133; laevis, 43; **raphani**, n. sp., cause of radish blackrot, 43

Aphid, on Douglas fir, 8; on silver fir, 9

Aphis, agents of transmission of dwarf disease, 639–646

Apothecia of Sclerotinia, two fungi on, 791–792

Apple, blotch, canker eradication, 46, 185–188; blotch fungus, details in life history of, 45–46; increasing prevalence of

Hypochnus rot of, 743–744; Pleospora rot of, 47; scab, control, 45; scab, infection studies, 45; "target canker" of, 735–738; thread blight of, 694; tree injection for fire blight, 51; trees, black walnut as a cause of the death of, 529–540; varieties, susceptibility to cedar rust, 541–543

Arachis hypogaea, pale dwarf disease of, 217–225

Armillaria mellea, cause of conifer root rot, 11

Arsenate of lead, for control of bacterial wilt and reduction of cucumber mosaic, 473–481

Ascochyta, blight of peas, relation of Mycosphaerella pinodes to, 44, 381–397; pisi, 381–397; pisi, description of, 381; species of, parasitic on pea, 381–397

Aspergillus niger, and scutellum rot of corn, 460

Avocado leaves, infected with Phyllachora gratissima, 153

BACH WALTER J., see WOLF, FREDERICK A.

Bacillus, amylovorus, control by tree injection, 51; amylovorus, flagella of, 405–406; carotovorus, cause of heart rot of celery, 663–667; tracheiphilus, control of cucumber wilt due to, 473–479

Bacteria, connected with target canker of apples and pears, 735–738

Bacterial, blight of pear, 611–627; blight of pear, symptoms of, 611–612; disease of Bowlesia, 507–512; halo spot of kudzu, 48

Bactericidal action of sodium silicofluoride, 50

Bacterium, campestre, serological differentiation from Bact. phaseoli, Bact. phaseoli sojense, and Bact. flaccumfaciens, 53–54; flaccumfaciens, serological and physiological studies of, 54; flaccumfaciens, serological differentiation from Bact. campestre, 53–54; phaseoli, serological and physiological studies of, 54; phaseoli, serological differentiation from Bact. campestre, 53–54; phaseoli sojense, correlation of virulence and acid agglutination in strains of, 49; phaseoli sojense, serological and physiological studies of, 54; phaseoli sojense, serological differentiation from Bact. campestre, 53–54; pruni, toxicity of sodium silicofluoride for, 50; **pueriae**, on kudzu, 48; solanacearum, on Hibiscus in Java, 25–27; tumefaciens, cause of sweet pea fasciation, 29–30; tumefaciens, strains of, 46–47

BAILEY, ALICE A., see RAMSEY, G. B.

BAIN, HENRY F., see STEVENS, NEIL E.

Barley, experiments with dusts controlling stripe disease of, 175–179; host of Aphanomyces euteiches, 133; loose smut, strains and host relationships, 42; seed, dehulling with sulphuric acid to induce infection with covered smut, 747–748

BARGER, W. R., see FAWCETT, H. S.

Bayer, dust, for control of barley stripe, 176–178

Bean rust (Uromyces appendiculatus), prevalence in Colombia, 152

Beef extract media versus beef infusion, 413–414

Beet, curly-top, virus of, causing curly-top of squash, 48–49; sugar, curly top of, in Argentine, 407–408; sugar, curly-top virus, cause of western yellow tomato blight, 39; sugar, ecological studies of curly-top of, 747; sugar, prevention of seedling diseases, 259–296

BENNETT, J. P., The treatment of lime-induced chlorosis in fruit trees, 745–746

Berberis, aristata, brachypoda, buxifolia, canadensis, chinensis, dictyophylla, leichlinii, lycium, pruinosa, thunbergii, vulgaris, nature of susceptibility, resistance, or immunity to Puccinia graminis, 98–111

Biography, of FITCH, HAROLD WAKEFIELD, 205–206; of LYMAN, GEORGE RICHARD, 355–361; of PIEMEISEL, FRANK J., 135–136; of SMITH, ERWIN F., 675–688

Biologic forms, see physiologic specialization

Birch, bronze borer, tree injection for, 51; serious disease of, 59

Blackberries, dwarf of, 629–648

Blackleg, of potato, seed-piece transmission, 155–160
Black-root, of radish, caused by **Aphanomyces raphani**, n. sp., 43
Black walnut (Juglans nigra) as a cause of the death of apple trees, 529–540
Blakeslea, distribution of 183–184; on Physalis angulata, 183; on Ipomoea batatas, 183; on tobacco leaves, 181–183; trispora, in culture of Botrytis rileyi, 181; trispora, occurrence of, in Dutch East Indies, 181–184
Blastophaga psenes L. (B. grossorum Grav.), the caprifying insect, transmitting Fusarium moniliforme var. fici, 199
Blight, Ascochyta, of peas, 44; bacterial, of pea, 611–627; chestnut, 11; citrus, in Florida, 58–59; Gloeosporium, of raspberry, 769–774; spur, of raspberry, differentiated from Gloeosporium blight, 769–770; thread, of citrus and pomaceous plants, 689–709; western yellow, of tomato, caused by beet curly-top virus, 39; western yellow, of tomato, in relation to environmental conditions, 746
BLISS, D. E., see MELHUS, I. E.
Blister rust, of white pine, 10
BLODGETT, F. M., A potato virus on peppers, 775–782; Tobacco mosaic on potatoes, 727–734
Blotch, apple, canker eradication, 46, 185–188; fungus, apple, details in life history, 45–46
Blue-mold rot, of citrus fruits, 47
Blue-stem (wilt), of raspberry, differentiated from Gloeosporium blight, 769–770
BOERNER, E. G., see TISDALE, W. H.
BOLLEY, H. L., The control of plant disease through seed certification—aided by test gardens and field crop inspections, 40
BONDE, REINER, Variation of strains of Alternaria solani from lesions on potato tubers, 56
Book review, Citrus diseases and their control, 203–204; Japanese textbook on plant pathology, 37; Manual of plant diseases, 341–342
Bordeaux mixture, and lead arsenate, for control of bacterial wilt and reduction of cucumber mosaic, 473–481; chemical hydrated lime for preparation of, 571–572
Bordeaux-oil emulsion, for citrus fruits, 47
Botanical congress, fifth international, 672–673
Botrytis rileyi, culture of, 181
Bowlesia septentrionalis, bacterial diseases, 507–512
BOWMAN, JOHN J., see FULTON, HARRY R.
BOYCE, J. S., Observations on forest pathology in Great Britain and Denmark, 1–18
BREWER, P. H., H. R. KRAYBILL, and MAX W. GARDNER, Purification of the virus of tomato mosaic, 744
BRIGGS, FRED N., Dehulling barley seed with sulfuric acid to induce infection with covered smut (Ustilago hordei), 747–748
BROWN, NELLIE A., Sweet pea fasciation, a form of crowngall, 29–30
Brown-patch of turf, fungicidal control of, 50–51
Brunchorstia destruens, cause of needle and twig blight of pines, 9
BRYAN, MARY K., Beef infusion versus beef extract media, 413–414; Flagella of Bacillus amylovorus, 405–406
Bud drop of gardenia, 671–672
BULLER, A. H. REGINALD, Cryptoporus volvatus and its relations with forest trees and insects, 53
Bunchy top, of abacá, or manila hemp, 255–257
Bunt, distribution of, 167–174
BUTLER, L. F., Increasing prevalence of Hypochnus rot of apples, 743–744; see ROSE, D. H.
Butt rot in Diospyros virginiana caused by Polyporus spraguei, 339–340

Cabbage, Chinese, Cercosporella leaf spot of, 42, 669–670; control of damping off, 503–504; Pythium ultimum on, 54–55
Caillieа glomerata (Forsk.) MacBride, 137
Calcium arsenate-gypsum, dust, for control of bacterial wilt of cucumbers, 473–481
Canker, apple blotch, eradication of, 46, 185–188; of apple blotch fungus, Phyllosticta solitaria E. and E., 185;

"Llaga" disease of coffee roots, 151; of Douglas fir, 4; of larch, 10; target, of apples and pears, 735–738
Capnodium brasilense Pat. and Maubl. causing sooty mold of coffee, 152
CARTER, WALTER, Ecological studies of curly-top of sugar beets, 747
Castilla cane susceptible to gomosis, 150
Cedar, leaf-blight, 11; rust, varietal susceptibility of apples to, 541–543
Celery, experiments, effect of mycorhiza, 746; heartrot, 663–667
Cell, effects of mosaic on content of, 57
Cells, of "mosaic diseased" and "dwarf" dahlias, x-bodies in, 52
Cenangium abietis, 9
Central Andes, plant disease notes from, 147–153
Cercospora (Helminthosporium) sacchari, cause of eye spot on sugar cane, 315, 317
Cercosporella (Cercospora), albo-maculans, cause of leaf spot of Chinese cabbage, 42; leaf spot of Chinese cabbage, 42, 669–670
Ceuthospora lunata, associated with storage rot of cranberries, 655
CHARDON, CARLOS E., and RAFAEL A. TORO, Plant disease notes from the central Andes, 147–153
Chemical, hydrated lime for the preparation of bordeaux mixture, 571–572; injury to tomatoes, 502–503; investigations of colored compounds in normal rust and mutants, 720–723; treatments of soil for control of damping-off fungi, 499–505
Chemicals, effect on germination of rust spores, 495–496
Chermes cooleyi, aphid on Douglas fir, 8
Cherry, Japanese, witches' broom of, 19–24; leaf spot, infection studies, 45
Chestnut blight, 11; tree injection for, 51
Chinese cabbage, Cercosporella leaf spot of, 42, 669–670
Chloroplasts, destruction of, in tomato mosaic, 365–367; physical and chemical properties of bodies associated with destruction of, in tomato mosaic, 367–372
Chlorosis, treatment of lime-induced, in fruit trees, 745–746
Choanephora, dichotoma Gandrup, 183; cucurbitarum, a true parasite on Amaranthus blitum, 183; infundibulifer (Cunn.) Sacc., 183; spp., 181–183
CHRISTENSEN, J. J. and E. C. STAKMAN, Susceptibility of wheat varieties and hybrids to wheat scab in Minnesota, 40–41; see STAKMAN, E. C.
Citrus, aurantium, infected by citrus scab, 153; "blight," wilt, or leaf-curl in Florida, 58–59; diseases and their control, book review, 203–204; fruit decay, temperature relations, 746–747; fruits, effect of spraying on keeping quality of, 47; plants, thread blight disease of, caused by Corticium koleroga, 689–709; scab, caused by Sphaceloma fawcetti, 153
CLAYTON, E. E., Effect of early spray and dust applications on later incidence of cucumber wilt and mosaic diseases, 473–481; Effect of seed treatment on growth, yield, and disease control in vegetable crops, 44
Clitocybe root-rot of trees and other woody plants in Florida, 56–57
Clover, leafhopper injury to, 58, 399–404
Coccomyces hiemalis, infection studies, 45
Coconut leaves, disease of, caused by Pestalozzia palmanum, 153
Coffee, root disease, 150–151
Collection, American type culture, 672
Colletotrichum, hibisci, on Hibiscus in Java, 25–27; lindemuthianum, 822
COLLEY, REGINALD H., and W. H. WESTON, Biography of GEORGE RICHARD LYMAN, 355–361
Color mutations of Puccinia graminis tritici, 711–725
Commercial tobaccos and cured leaf as a source of mosaic disease in tobacco, 49, 513–522
Congress, fifth international botanical, 672–673
Conidia, living, photographing on agar, 195–198
Coniferous seed-beds, damping-off control in, 51–52
Coniothyrium spp., connected with target canker of apples and pears, 736

Control, of barley stripe with chemical dusts, 175–179; of brown-patch of turf by fungicides, 50–51; of common scab of potato with sulphur, 297–314; of cotton root rot caused by Fusarium moniliforme, 235–237; of cotton wilt, 765; of damping off fungi, by chemical treatment of soil, 499–505; of damping-off in coniferous seed-beds, 51–52; of dwarf of blackberries, 646–647; of fungous diseases and insect pests by tree injection, 51; of internal rot of caprified figs, 199–200; of millet smut, 739–741; of oat smut in Idaho, dust treatment for, 569–570; of bacterial blight of pea, 625–626; of plant disease through seed certification—aided by test gardens and field crop inspections, 40; of Rhizoctonia of potatoes in Idaho, seed treatment, 793–814; of sand burn of pecan seedlings, 660; of sorghum kernel smut and millet smut by sulphur and copper carbonate dusts, 52; of thread blight, of citrus and pomaceous fruits, 706–707

COOK, MELVILLE T., Some effects of mosaic on the content of the cell, 57

COONS, G. H. and DEWEY STEWART, Prevention of seedling diseases of sugar beets, 259–296

Copper carbonate, for control of bunt, 175; for control of damping off of tomatoes, 499–505; for control of millet smut, 740; for control of oat smut, 569–570; for control of sorghum kernel smut and millet smut, 52

Copper, correction of unproductive muck by the addition of, 49–50; dusts, used on lettuce and onions on muck soils, 49–50

Corn, downy mildew, 57; fungous flora of nodes of, 41; mosaic, of Hawaii, distinct from sugar cane mosaic, 41; nature of resistance to Puccinia sorghi, 815–825; Ophiobolus heterostrophus on, 414; permeability of seed coat to mercury compounds, 51; resistance to smut, 42; rust, see Puccinia sorghi L.; scutellum rot, 449–471; smut, see Ustilago zeae; study of fungous flora of nodal tissues of, 563–568

Correlation of virulence and acid agglutination in strains of Bacterium phaseoli sojense, 49

Corrosive sublimate, see mercuric chloride

Corticum, koleroga, cause of thread blight on citrus and pomaceous fruits, 689–709; koleroga, morphology of, 700–702; koleroga, pathogenicity of, 699–700; pathological anatomy of affected plants, 702; portentosum, 137; stevensii, 705; vagum var. solani, effect on potatoes, 811

Cotton, root diseases caused by Fusarium moniliforme Sheld., 227–238; seed, a modified method of delinting, 189–193; wilt of, 749–767

Covered smut, dehulling barley seed with sulphuric acid to induce infection with, 747–748; of oats, effects of soil moisture, soil temperature, and dehulling on, 31–36

CRAIGIE, J. H., see MELANDER, L. W.

Cranberries, storage rots of, 649–655

Cronartium ribicola, 10

Crowngall, producing fasciation of sweet pea, 29–30; strains of causal organism, 46–47

Crown rust, of oats, physiologic specialization in, 783–790

Cryptoporus volvatus and its relations with forest trees and insects, 53

Cucumber, beetle, striped (Diabrotica vittata), control, 473–481; wilt and mosaic, control by fungicides, 473–481; variety of, resistance to mosaic, 48

Culture, American type, collection, 672

Cultures, fungous, relation of storage temperature to lag in growth of, 55

Cured leaf, and commercial tobaccos, as source of tobacco mosaic, 49, 513–522

"Curly top," of abacá, or Manila hemp, 255

Curly-top, of sugar beets, ecological studies of, 747; of sugar beet in Argentine, 407–408; of squash, 48–49; virus, of sugar beet, cause of western yellow tomato blight, 39

Currant, blister rust on, 10

Cytological study of Puccinia coronata on Banner and Cowra oats, 748

Dahlias, "mosaic diseased" and "dwarfed," x-bodies in cells of, 52
Damping-off, control in coniferous seedbeds, 51–52; control of, by soil treatment with chemicals, 500–503; Pythium debaryanum the cause of, 54–55
Dasyscypha calycina, cause of larch canker, 10
DAVIS, W. H., Cercosporella leaf spot of Chinese cabbage, 42; Notes on the Cercosporella leaf spot of Chinese cabbage in Massachusetts, 669–670
Decay, citrus fruit, temperature relations, 746–747
Decays, fruit, of Feijoa, 745
Dehulling, of barley seed with sulphuric acid to induce infection with covered smut, 747–748; of oats, effect on smut infection, 31–36
Delinting cotton seed, a modified method of, with sulphuric acid, 189–193
DEMAREE, J. B., Sand burn of pecan seedlings, 657–661
Dematophora sp., conidial stage of Rosellinia fungus, 151
Denmark, observations on forest pathology in, 1–18
Desmanthus, 137
Diabrotica vittata, striped cucumber beetle, control, 473–481
Diaporthe citri, 47
Dichrostachys nutans, a weed tree, in Cuba, with remarks on its pathology, 137–146
DICKSON, JAMES G., see LEUKEL, R. W.
Differential hosts of Puccinia coronata avenae, 787
Differentiation, serological, of Bacterium campestre from Bact. phaseoli, Bact. phaseoli sojense, and Bact. flaccumfaciens, 53–54
Diospyros virginiana, butt rot of, 339–340
Diplodia natalensis, 47
Disease, bacterial, of Bowlesia, 507–512; "Llaga" (canker), on coffee roots, 151; nematode, of sweet potato, 549–554; of avocado leaves, caused by Phyllachora gratissima, 153; of birches, 59; of coconut leaves caused by Pestalozzia palmanum Cooke, 153; of coffee root, 150–151; of larkspur, caused by Sclerotium delphinii, 242–244; plants, notes from the Central Andes, 147–153; scutellum rot of corn, 499–471; seedling, of sugar beets, 259–263
Dissemination, of bacterial blight of pea, 611–627
DODGE, B. O., A Gloeosporium blight of raspberry, 769–774
Dothichiza populea and its mode of infection, 545–547
Douglas fir, aphid, 8; canker of, 4; needle cast of, 7
Downy mildew, Sclerospora graminicola, a study of, 57
DRECHSLER, CHARLES, A peculiar type of Pythium, 55; An emendation of the description of Ophiobolus heterostrophus, 414; Pythium ultimum and P. debaryanum, 54–55
Dreyfusia nuesslini, aphid on silver fir, 9
Drop, bud, of gardenia, 671–672
DUFF, G. H., and CATHERINE GRAHAM WELCH, Sulfur as a control agent for common scab, 297–314
Du Pont, dusts, for control of barley stripe, 176–178; dusts, for control of oat smut, 570; dusts, toxic effect on fungous spores, 317, 218; No. 15, dust, for control of Rhizoctonia of potatoes, 795, 814
Dust, effect of early applications, on cucumber wilt and mosaic diseases, 473–481; fungicides, method of testing toxicity of, 315–319; treatment for control of oat smut in Idaho, 569–570
Dusts, experiments with, for controlling stripe disease of barley, 175–179; sulphur and copper carbonate for control of sorghum and millet smuts, 52
Dwarf, of blackberries, 629–648; pale, of peanut, 217–225; yellow, disease of potatoes, on peppers, 775–782
Dwarfing of peanuts, differences between seed lots, 220–221; relation of physical factors to, 222–224; relation of seed vigor to, 222
DYKSTRA, T. P., see MCKAY, M. B.

ECKERSON, SOPHIA H., and H. R. KRAYBILL, Separation of fern leaf from mottling in tomato mosaic, 57–58
Ecological studies of curly-top of sugar beets, 747

Effect, of a certain strain of tobacco mosaic on the yield and quality of burley tobacco, 523–527; of different hosts upon the sporangia of some Phytophthoras, 483–490; of early spray and dust applications on later incidence of cucumber wilt and mosaic diseases, 473–481

Effects of fungous extracts upon the initiation and growth of the perithecia of Venturia inaequalis (Cke.) Wint. in pure culture, 835–836

Egypt, distribution of cotton wilt in, 750–752

Elachiptera costata, agent of inoculation of celery heart rot, 663–670

Elm, disease, 13; European scale, tree injection for, 51

ELMER, O. H., A mosaic-resistant variety of cucumbers, 48

Emendation of the description of Ophiobolus heterostrophus, 414

Empoasca, fabae, leafhopper causing yellowing of alfalfa, 39; fabae on clover, 399–404; mali, leafhopper injuring clover, 58

End rot, of cranberries in storage, 652

Endothia parasitica, 11; tree injection for, 51

Environmental conditions in relation to western yellow tomato blight, 746

Enzymes, of Pythiacystis citrophthora, 747

Eradication, of apple blotch canker, 46, 185–188

Erysiphe scandens, 705

Eutettix tenella, carrier of beet curly-top virus, 39; in connection with western yellow tomato blight, 746; leafhopper carrying squash curly-top virus, 48–49

Exoascus cerasi, cause of witches' broom on Japanese cherry trees, 19–24; hosts of, 22

Exobasidium, **parvifolii** Hotson, n. sp., 214–215; spp., discussion of, 207–208

Extract, beef, versus beef infusion media, 413–414

Extracts, effects of fungous, upon development of perithecia, 835–836

Eye spot of sugar cane, 315

EZEKIEL, WALTER N., Two fungi on Sclerotinia apothecia, 791–792

Factors, genetic, for inheritance of rust reaction and growth habit in wheat, 573–609

FAHMY, TEWFIK, Fusarium disease (wilt) of cotton and its control, 749–767

FARIS, J. A., Zonate foot rot of sugar cane, 83–94; and R. V. ALLISON, Sugar cane root disease in Cuba: a progress report upon the root disease situation in 1925, 61–82

Fasciation, of sweet pea, form of crown-gall, 29–30

FAWCETT, G. L., Curly top of sugar beet in the Argentine, 407–408

FAWCETT, H. S., and W. R. BARGER, Relation of temperature to growth of Penicillium italicum and P. digitatum and to citrus fruit decay produced by these fungi, 746–747; and H. ATHERTON LEE, Citrus diseases and their control, textbook, review of, 203–204

Feijoa sellowiana, fruit decays, 745

FELIX, E. L., Correction of unproductive muck by the addition of copper, 49–50

Fern-leaf, separation from mottling in tomato mosaic, 57–58

Feterita, loose kernal smut on, 338–339

Fifth international botanical congress, 672–673

Fig rust, Kuehneola fici, description of, 152

Figs, caprified, control of internal rot of, 199–200

Fir, Douglas, aphid, 8; Douglas, canker of, 4; Douglas, needle cast of, 7; Douglas, root rot of, 11; silver, aphid on, 9; silver, root rot of, 11; twig and needle, blight of, 9

FITCH, HAROLD WAKEFIELD, biography of, 205–206; in memoriam, 353

Fixing, of corn rust material, 816

Flagella of Bacillus amylovorus, 405–406

FLINT, W. P., see HOLLOWELL, E. A.

Flora, study of fungous, of nodal tissues of corn plant, 563–568

FOLSOM, DONALD, Uniformity of nomenclature for the viroses of Solanum tuberosum, 161–165

Fomes annosus, cause of conifer root rot, 11, 14; annosus, strains of, 12; sp., pos-

sible cause of zonate foot rot of sugar cane, 92–94
Footrot, zonate of sugar cane, 83–94
Forest, parasites, strains of, 12; pathology, observations on, in Great Britain and Denmark, 1–18; trees, relation of Cryptoporus volvatus to, 53
Formaldehyde, for control of oat smut, 569–570; for control of Rhizoctonia of potatoes, 795–814
Forms, see physiologic specialization
FRASER, W. P., see RUTTLE, MABEL L.
FROMME, F. D., S. A. WINGARD, and C. N. PRIODE, Ringspot of tobacco; an infectious disease of unknown cause, 321–328
Fruit, citrus, decay, temperature relations, 746–747; decays of Feijoa, 745; trees, treatment of lime-induced chlorosis in, 745–746; citrus, effect of spraying on keeping quality of, 47
FULTON, HARRY R., and JOHN J. BOWMAN, The effect of spraying with fungicides on the keeping quality of Florida citrus fruits, 47
Fungicidal control of brown-patch of turf, 50–51; value of oil sprays, 201–202
Fungicides, control of cucumber diseases by, 473–481; control of damping off by, 499–505; dust, a method for testing, toxicity of, 315–319; effect of spraying with, on keeping quality of citrus fruits, 47; sulphur and copper carbonate dusts for sorghum and millet smut control, 52
Fungous, cultures, relation of storage temperature to lag in growth of, 55; diseases, control of, by tree injection, 51; extracts, effects of, upon the initiation and growth of the perithecia of Venturia inaequalis (Cke.) Wint. in pure culture, 835–836; flora, of nodal tissues of corn plant, 563–568; spores, testing toxicity of dust fungicide to, 315–319
Furfural, for control of Rhizoctonia of potatoes, 795
Fusarium, cubense, in connection with heart rot of abacá, or Manila hemp, 255; culmorum (W. G. Sm.) Sacc., living conidia, photographed, 197; disease of cotton and its control, 749–767; eumartii Carp., living conidia, photographed, 197; martii App. u. Wr., living conidia, photographed, 197; moniliforme Sheld. causing internal rot of figs, 199; moniliforme Sheld., on cotton roots, 227–238; moniliforme Sheld. var. fici Caldis, 199; niveum, infection studies, 47–48; oxysporum Schl., living conidia, photographed, 197; solani (Mart. pr. p.) App. u. Wr., living conidia, photographed, 197; sp., and scutellum rot of corn, 460; sp. cause of wilt disease of alfalfa, 337–338; sp., on Sclerotinia americana, 791–792; spp. in corn nodes, 41; vasinfectum Atk., living conidia, photographed, 197; vasinfectum, relation to Egyptian and Indian Fusarium, 761–764; wilt of sweet potatoes, 42–43
Fusicoccum putrefaciens, cause of end rot of cranberries in storage, 652

Ganoderma pulverulentum Murr., 142; associated with Ustulina zonata (Lev.) Sacc., 142; causing decay of heartwood, 145
Gardenia bud drop, report of, 671–672
GARDNER, MAX W., Apple blotch canker eradication, 46, 185–188; see BREWER, P. H.
Genetics, of growth habit and rust reaction in wheat crosses, 573–609
Germination, of rust spores, 491–498; of rust spore, of color mutants, 713–715
Germisan, for control of barley stripe, 175
Gibberella saubinetii, and scutellum rot of corn, 460
GILBERT, A. H., Net necrosis of potato, 555–561
GILLIGAN, G. M., see HOLLAND, E. B.
Gloeosporium, blight of raspberry, 769–774; cingulatum, on raspberry, 771; sp., associated with storage rot of cranberries, 655; venetum, on raspberry, 772–773; cingulata, on raspberry, 771
GOLDSTEIN, BESSIE, The x-bodies in the cells of "mosaic diseased" and "dwarfed" dahlias, 52
"Gomosis" of sugar cane, description of, 148–150
Gossyparia spuria, European elm scale, 51
GRANOVSKY, A. A., see JONES, FRED R.

Grapefruit, thread blight of, 693
GRAVATT, ANNIE RATHBUN, see RATHBUN-GRAVATT
Great Britain, observations on forest pathology in, 1–18
GRIFFITHS, MARION A., Smut resistance in corn, 42; see TISDALE, W. H.
Growth habit, in crosses between spring and winter wheats, 578–595
Guignardia vaccinii, associated with storage rot of cranberries, 652
Gymnosporangium juniperi-virginianae, varietal susceptibility of apples to, 541–543

Halo spot, bacterial, of kudzu, 48
HANSEN, H. N., Control of internal rot of caprified figs, 199–200
HARMON, T. CARTER, see MONTEITH, JOHN JR.
HARTLEY, CARL, Notes on Hibiscus diseases in west Java, 25–27; Pale dwarf disease of peanut, Arachis hypogaea, 217–225
HASKELL, R. J., Report of 18th annual meeting of the American phytopathological society, 343–352
HEALD, F. D., Manual of plant diseases, book review, 341–342
Heart rot of abacá, or Manila hemp, 255–257
Heart rot, of celery, 663–667
HEDGECOCK, GEORGE G., Dothichiza populea and its mode of infection, 545–547
HEDGES, FLORENCE, Bacterial halo spot of kudzu, 48
Helminthosporium (Cercospora) sacchari, cause of eye spot on sugar cane, 315, 317; spp., in corn nodes, 41
Heterodera radicicola, in connection with bunchy top of abacá, or Manila hemp, 255; radicicola, on Hibiscus in Java, 26
Heterothallism in Ustilago zeae, 827–834
Hevea, diseases of, 143
Hibiscus diseases in west Java, 25–27
HIGGINS, B. B., Physiology and parasitism of Sclerotium rolfsii, 53, 417–448
H-ion concentration, relation to growth of Sclerotium rolfsii, 420–428
Histological study, of Puccinia sorghi, 815–825; pathological, of bacterial blight of pea, 611–627
HOLLAND, E. B., and G. M. GILLIGAN, Chemical hydrated lime for the preparation of bordeaux mixture, 571–572
HOLLOWELL, E. A., JOHN MONTEITH, JR., and W. P. FLINT, Leafhopper injury to clover, 58; Leafhopper injury to clover, 399–404
HORNE, W. T., Notes on fruit decays of the feijoa (Feijoa sellowiana Berg), 745
Host, morphology, of wheat stems and seedling leaves in relation to rust reaction, 602–603; plants, see parasitism, also differential hosts; range, see parasitism; tissue, relation of Sclerotium rolfsii to, 442–443
HOTSON, J. W., New species of Exobasidium, 207–216
HUNGERFORD, C. W., see RAEDER, J. M.
Hydrated lime, chemical, for preparation of bordeaux mixture, 571–572
Hypochnopsis (ochroleuca) ochroleucus, 705
Hypochnus rot of apples, increasing prevalence of, 743–744

Ikeno, S., Review of Suematu's textbook on plant pathology in Romanized Japanese, 37
Increasing prevalence of Hypochnus rot of apples, 743–744
Infection, Dothichiza populea and its mode of, 545–547
Infection studies, watermelon wilt, 47–48
Infections, twig, by cedar rust, susceptibility of apple varieties to, 541–543
Infusion, beef, versus beef extract media, 413–414
Inheritance, of growth habit in crosses between spring and winter wheats, 573–609; of rust reaction in crosses between Marquis, Kota, and Kanred wheats, 573–609
Injection, tree, for control of fungous diseases and insect pests, 51
Injury, leafhopper, to clover, 58; 399–404
In memoriam, 353
Insect pests, control of, by tree injection, 51; relation of Cryptoporus volvatus to, 53; relation of, to the development of heart rot of celery, 663–667

Internal rot of caprified figs, control of, 199–200
International, fifth botanical congress, 672–673
Investigations on citrus "blight," wilt or leafcurl in Florida, 58–59
Ipomea batatas, infected by Blakeslea, 183

Japanese cherry, witches' broom of, 19–24; larch, canker of, 6
Java, Hibiscus diseases in, 25–27
JOCHEMS, S. C. J., The occurrence of Blakeslea trispora Thaxter in the Dutch East Indies, 181–184
JOHNSON, A. G., see LEUKEL, R. W.
JOHNSON, E. M., see VALLEAU, W. D.
JOHNSON, THORVALDUR, see NEWTON MARGARET
JOHNSTON, C. O., Effects of soil moisture and temperature and of dehulling on the infection of oats by loose and covered smuts, 31–36; see MELCHERS, L. E.
JONES, LEON K., The relation of Mycosphaerella pinodes to Ascochyta blight of peas, 44
JONES, FRED R., and A. A. GRANOVSKY, Yellowing of alfalfa caused by leafhoppers, 39
Juglans nigra L., as a cause of the death of apple trees, 529–540
Jute, diseases of in Java, 25

Karasch, compounds, for control of barley stripe, 176–178
Keithia thujina, cause of cedar leaf-blight, 11
KEITT, G. W., Studies of apple scab and cherry leaf spot infection under controlled conditions, 45; and E. E. WILSON, A possible reorientation of aims and methods for apple scab control, 45
KENDRICK, JAMES B., Radish black-root caused by **Aphanomyces raphani**, n. sp., 43
Kernel smut, loose, on Feterita, 338–339; of sorghum, control, 52
Key for identification of physiologic forms of Puccinia coronata avenae, 787
KLOTZ, L. JOS., The enzymes of Pythiocystis citrophthora Sm. and Sm., 747
KOHL, EDWIN J., Details in the life history of the apple blotch fungus, 45–46
KOEHLER, BENJAMIN, Studies on the scutellum rot disease of corn, 449–471
KRAYBILL, H. R., see BREWER, P. H.; see ECKERSON, SOPHIA H.
Kudzu, bacterial halo spot of, 48
Kuehneola fici, cause of fig rust, 152
KUNKEL, L. O., The corn mosaic of Hawaii distinct from sugar cane mosaic, 41

Lag in growth of fungous cultures, relation of storage temperature to, 55
Larch, canker, 10; Japanese, canker of, 6; needle cast of, 10
Larkspur, Sclerotium disease of, 239–245
Lathyrus, latifolius, odoratus, sp., host of Aphanomyces euteiches, 133
LEACH, J. G., The nature of seed-piece transmission of potato blackleg, 155–160; The relation of insects and weather to the development of heart rot of celery, 663–667
Lead arsenate, for control of bacterial wilt and reduction of cucumber mosaic, 473–481
Leaf blight, of cedar, 11
Leaf-curl, citrus, in Florida, 58–59
Leafhopper, injury to clover, 58; injury to clover, 399–404; beet, carriers of squash curly-top, 48–49; cause of yellowing of alfalfa, 39
Leaf, rust, effect on seed production of wheat, 40; spot, Cercosporella, of Chinese cabbage, 42, 669–670; spot of coffee, 151; spot of cocoanuts, 153; spot of tomatoes, 153
Leaves, detached, multiplication of tobacco mosaic virus in, 58
LE CLERG, E. L., Leaf temperature in relation to tip burn of lettuce, 44–45
LEE, H. ATHERTON, and J. P. MARTIN, A method for testing in vitro the toxicity of dust fungicides to fungous spores, 315–319; see HOWARD S. FAWCETT
LEIGHTY, C. E., see TISDALE, W. H.
Lemons, Pleospora rot, 47
LEONIAN, LEON H., The effect of different hosts upon some species of Phytophthora, 56; The effect of different hosts upon

the sporangia of some Phytophthoras, 483–490
Leptosphaeria sacchari, cause of ring spot of sugar cane, 153
Lettuce, on muck soils, use of copper on, 49–50; tip burn, relation of leaf temperature to, 44–45
LEUKEL, R. W., JAMES G. DICKSON, and A. G. JOHNSON, Experiments with dusts for controlling stripe disease of barley, 175–179
LEWIS, I. M., and ELIZABETH WATSON, A bacterial disease of Bowlesia, 507–512
Light, effect on germination of rusts, 493–494
Lime, chemical hydrated, for preparation of bordeaux mixture, 571–572; induced chlorosis in fruit trees, treatment of, 745–746
LINFORD, M. B., Additional hosts of Aphanomyces euteiches, the pea rootrot fungus, 133–134
LINFORD, MAURICE B., and RODERICK SPRAGUE, Species of Ascochyta parasitic on pea, 381–397
LINK, GEO. K. K., and C. G. SHARP, Serological differentiation of Bacterium campestre from Bact. phaseoli, Bact. phaseoli sojense, and Bact. flaccumfaciens, 53–54
"Llaga" disease found at the "Amparo" coffee plantation, Colombia, 151
Loose kernel smut on Feterita, 338–339
Loose smut, of oats, effects of soil moisture, soil temperature, and dehulling on, 31–36
LYMAN, GEORGE RICHARD, biography of, 355–361; in memoriam, 353
Lycopersicon esculentum Mill. virosis 2, synonymy, 164

MAINS, E. B., The effect of leaf rust, Puccinia triticina, on the seed production of wheat, 40
Maize, see corn
MANEVAL, W. E., Further germination tests with teliospores of rusts, 491–498
Manila hemp, bunchy top and heart rot of, 255–257
Manual of plant diseases, book review, 341–342
Marabu, diseases of, 142; eradication of, 140; importance as a pest, 138; roots of, 139; used for charcoal, 142
MARTIN, J. H., and G. T. RATLIFFE, Loose kernel smut on Feterita, 338–339
MARTIN, J. P., see LEE, H. ATHERTON
MASSEY, A. B., Book review, Manual of plant diseases, 341–342; Biography of HAROLD WAKEFIELD FITCH, 205–206
MCKAY, M. B., and T. P. DYKSTRA, Curly-top of squash, 48; Sugar beet curly-top virus, the cause of western yellow tomato blight, 39
MCWHORTER, FRANK P., Fungicidal value of oil sprays, 201–202
Media, beef infusion versus beef extract, 413–414; growth of Sclerotium rolfsii on, 418–419
Medicago sativa, host of Aphanomyces euteiches, 133
Melampsora lini, 822
Melanconium iliau, association with root disease of sugar cane, 63–67; iliau, association with zonate footrot of sugar cane, 91; sacchari, association with root disease of sugar cane, 63–68; sacchari Massee, a rind disease of sugar cane, caused by, 152; sacchari, association with zonate footrot of sugar cane, 91
MELANDER, L. W., and J. H. CRAIGIE, Nature of resistance of Berberis spp. to Puccinia graminis, 95–114
MELCHERS, L. E., Studies on the control of millet smut, 739–741; and C. O. JOHNSTON, Sulphur and copper carbonate dusts as efficient fungicides for the control of sorghum kernel smut and millet smut, 52
MELHUS, I. E., FRANK VAN HALTERN, and D. E. BLISS, A study of the downy mildew, Sclerospora graminicola (Sacc.) Schroet., 57
Melilotus alba, host of Aphanomyces euteiches, 133
Mercuric, chloride, for control of damping off of tomatoes, 499–505; chloride, for control of oat smut, 569–570; chloride, for control of Rhizoctonia of potatoes, 795–814; fungicides for brown-patch control, 50–51

Mercury "C," for control of barley stripe, 176–178; compounds for control of Rhizoctonia in cabbage and tomato plantings, 499–505; compounds, injurious to tomatoes in concentrations not harmful to cabbage, 499–505; compounds, permeability of corn seed coat to, 51
Meria laricis, cause of needle cast of larch, 10
Metabolic products, of Sclerotium rolfsii, 428–434
Method, for testing in vitro the toxicity of dust fungicides to fungous spores, 315–319; of delinting cotton seed with sulphuric acid, 189–193; of studying nature of resistance of Berberis spp. to Puccinia graminis, 97–98
Micrococcus ulmi, cause of elm disease, 13
Mildew, downy, Sclerospora graminicola, a study of, 57; powdery, of muskmelon, 414–415; rose, caused by Sphaerotheca parmosa, 201
Millet smut, control studies, 739–741; control with sulphur and copper carbonate dusts, 52
Mimosa, 137
Mimosidiae, 137
Moisture, soil, effect on smut infection of oats, 31–36
MONTEITH, JOHN JR., and T. CARTER HARMON, Fungicidal control of brown-patch of turf, 50–51; see HOLLOWELL, E. A.
Morphology, effect of hosts on, of Phytophthora sporangia, 483–490; of wheat stems and seedling leaves in relation to rust reaction, 602–603
"Mosaic bodies" of tomato leaf, 372–376
Mosaic, cucumber, control, 479–480; dahlias, x-bodies in cells of, 52; destruction of chloroplasts in tomato, 363–379; disease in tobacco, commercial tobaccos and cured leaf as source of, 513–522; disease in winter wheat, certain factors influencing development of, 41; effects on content of the cell, 57; in "natural leaf" chewing tobacco, 517; of corn in Hawaii, distinct from sugar cane mosaic, 41; of grasses, cane, and corn, 152; of raspberries, 46; of tobacco, 152; tissue, histological changes in, 366–367; tobacco, effect of strain of, on yield and quality of burley tobacco, 523–527; tobacco, from commercial and cured leaf tobaccos, 49; tobacco, multiplication of virus in detached leaves, 58; tobacco, on potatoes, 727–734; tomato, purification of virus, 744; tomato, separation of fern leaf from mottling in, 57–58; suggested uniformity of nomenclature, 161–165; variety of cucumbers resistant to, 48; virus, tobacco, effect of age on, 515–517
Mottling, separation of fern leaf from, in tomato mosaic, 57–58
Muck, unproductive, correction of with copper, 49–50
Mucor sp., and scutellum rot of corn, 460
Multiplication of the virus of tobacco mosaic in detached leaves, 58
Muskmelon, powdery mildew of, 414–415
Mutants of Puccinia graminis tritici, color, 711–725
Mycorrhiza, celery experiments, 746
Mycosphaerella pinodes, relation to Ascochyta blight of peas, 44, 381–397; pinodes, description of, 390–391

Nailhead spot of tomatoes, development during transit, 43–44
Nature of resistance in Zea mays to Puccinia sorghi, 815–825
"Natural leaf" chewing tobacco, mosaic in, 517
Necrosis, net, of potato, 555–561
Nectria, causing birch disease, 59
Needle and twig blight, of firs, 9; of pines, 9
Needle cast, of Douglas fir, 7; of larch, 10
Nematode disease of sweet potato, 549–554
Nematodes, on Hibiscus in Java, 26
Net necrosis of potato, 555–561
NEWTON, MARGARET, and THORVALDUR JOHNSON, Color mutations in Puccinia graminis tritici, 711–725
Nicandra physalodes, potato virus on, 775–781
Nicotiana tabacum L. virosis 5, synonymy, 164

Nodal tissues of corn plant, a study of the fungous flora of, 563–568
NOLLA, J. A. B., A new Alternaria disease of onions (Allium cepa L.), 115–132
Nomenclature, uniformity of, for viroses of Solanum tuberosum, 161–165
Notes, on the Cercosporella leaf spot of Chinese cabbage in Massachusetts, 669–670; phytopathological, summer meeting, 836
Nuclear condition of color mutants of Puccinia graminis tritici, 719–720

Oats, effects of soil moisture, soil temperature, and dehulling on smut infection of, 31–36; host of Aphanomyces euteiches, 133; smut, dust treatment for control of, in Idaho, 569–570; Banner and Cowra, cytological study of Puccinia coronata on, 748
OCFEMIA, GERARDO OFFIMARIA, Second progress report on bunchy-top of abacá, or manila hemp, 255–257
Odostemon, aquifolium, nature of resistance to Puccinia graminis, 98–111; repens, nature of immunity to Puccinia graminis, 98–111; swaseyi, nature of susceptibility to Puccinia graminis, 98–111
Oil sprays, fungicidal value, 201–202
Omphalia flavida Maubl. et Rang., 151
Onions, on muck soils, use of copper on, 49–50
Ophiobolus heterostrophus, emendation of description of, 414
Orange, thread blight of, 693
ORTON, C. R., The permeability of the seed coat of corn to mercury compounds, 51
Overwintering, of bacterial blight of pea, 611–627
Oxalic acid, production by Sclerotium rolfsii, 428–434; toxicity of, 434–441

Pacific Division, American Phytopathological Society, abstracts of, 745–748
Pale dwarf disease of peanut, 217–225
Parasite, use of term, 409–411
Parasites, forest, strains of, 12–13
Parasitic strains, see physiologic specialization
Parasitism (host plants, host range, pathogenicity), color mutants, Puccinia graminis tritici, 713; cotton wilt, susceptibility of varieties, 757–758; effects of hosts on some species of Phytophthora, 56; range of Pseudomonas pisi, 623–625; oat varieties, susceptibility to Puccinia coronata avenae, 787; of Sclerotium rolfsii, 53; physiology and, of Sclerotium rolfsii, 417–448; varietal susceptibility to dwarf of blackberries, 631–633
PARSON, H. E., Physiologic specialization in Puccinia coronata avenae, 783–790
PATEL, M. R., Strains of Pseudomonas tumefaciens, 46–47
Pathogenicity, see Parasitism
Pathology, forest, observations on, in Great Britain and Denmark, 1–18
Pea, bacterial blight of, 611–627; root rot of, 133–134; species of Ascochyta parasitic on, 381–397; sweet, fasciation, form of crowngall, 29–30
Peach, tissues not injured by sodium silicofluoride, 50
Peanut, pale dwarf disease of, 217–225
Pear, "target canker" of, 735–738; thread blight of, 694
Peas, Ascochyta blight of, 44
Pecan, seedlings, sand burn of, 657–661; thread blight of, 696
Pellicularia koleroga, 705
Penicillium, digitatum, 47; digitatum, temperature relations, 746–747; effect of pure culture on development of the perithecia of Venturia inaequalis, 835–836; italicum, 47; sp., and scutellum rot of corn, 460
Peppers, potato virus on, 775–782
Peregrinus maidis, corn leafhopper, unable to transmit sugar cane mosaic, 41
Perithecia, of Venturia inaequalis, effects of fungous extracts in pure culture, 835–836
Permeability of seed coat of corn to mercury compounds, 51
Pestalozzia palmanum Cooke causing disease of cocoanut leaves, 153
PIEMEISEL, FRANK J., biography of, 135–136

Phoma, abietina, 4; betae, cause of seedling disease of sugar beet, 263; pithya, 5

Phomopsis citri, 47; pseudotsugae, cause of canker of Douglas fir, 4; cause of canker of Japanese larch, 6; pseudotsugae, strains of, 12; sp. associated with storage rot of cranberries, 652

Photographing living conidia on agar, 195–198

Phthora vastatrix, a Pyrenomycete, cause of coffee root disease, 150

Phyllachora gratissima on avocado leaves, 153

Physalis angulata, infected by Blakeslea, 183

Phyllosticta solitaria E. and E., the apple blotch fungus, 185

Physalospora, rhodina, relation of temperature to, 55; malorum, connected with target canker of apples and pears, 736

Physical factors, relation to pale dwarf of peanuts, 222–224

Physiologic, specialization in Puccinia coronata avenae, 783–790; specialization, in Tilletia tritici and T. levis, 247–253; specialization, in Puccinia sorghi, 815–825; specialization, of stem rust, factors for inheritance of reaction to physiologic forms in wheat, 595–601; strains, smooth and rough, of Bacterium phaseoli sojense, 49; strains, of Alternaria solani from potato tubers, 56; strains of forest parasites, 12–13; strains, of Pseudomonas tumefaciens, 46–47; strains of Ustilago nuda, 42

Physiological studies of Bacterium phaseoli, Bact. phaseoli sojense, and Bact. flaccumfaciens, 54

Physiology, of Sclerotium rolfsii, 53; and parasitism of Sclerotium rolfsii, 417–448

Phytomonas bowlesii, cultural and physiological relations, 508–511; n. sp., description, 511–512; disease of Bowlesia due to, 508–511

Phytopathological Society, see American Phytopathological Society

Phytophthora, effect of different hosts upon some species of, 56; nicotianae, causes same symptoms as Choanephora, 183; spp., damping off due to, 500; spp., effect of different hosts on morphology, 483–490

Pine, Corsican, root rot of, 11; white, blister rust, 10; needle and twig blight of, 9

Pisum sativum, host of Aphanomyces euteiches, 133; species of Ascochyta on, 381–397

Plant, disease control by seed certification, 40; disease notes from the central Andes, 147–153; pathology, review of Japanese text-book, 37

Pleospora herbarum, var. citrorum, 47; rot of lemons and apples, 47; sp., relation of temperature to, 55

Polyporus opacus Berk. and Mont., species of, 142; spraguei, cause of butt rot of Diospyros virginiana, 339–340

Polystictus pinsitus, 137

Pomaceous plants, thread blight disease of, caused by Corticium koleroga, 689–709

Pomegranate, thread blight of, 696

POOLE, R. F., The Fusarium wilt of sweet potatoes on infested soils, 42; and ROBERT SCHMIDT, The nematode disease of sweet potatoes, 549–554

Populus spp. strains of Taphrina aurea on, 12

PORTER, D. R., Watermelon wilt infection studies, 47–48

PORTER, C. L., A study of the fungous flora of nodal tissues of the corn plant, 563–568; the fungous flora of the nodes of corn, 41

Potato, blackleg, nature of seed-piece transmission of, 155–160; net necrosis, 555–561; seed treatment control of Rhizoctonia, in Idaho, 793–814; sulfur as a control agent for common scab of, 297–314; sweet, nematode disease of, 549–554; tubers, strains of Alternaria solani from, 56; uniformity of nomenclature for viroses of, 161–165; virus on peppers, 775–782; Bliss Triumph, inoculated with tobacco mosaic virus, 729–730; Green Mountain, inoculated with tobacco mosaic virus, 728–729; inoculated with tobacco mosaic virus, 727–734

Powdery mildew of muskmelon, 414–415
Preliminary report on gardenia bud drop, 671–672; studies on witches' broom of strawberry, 329–335
Preparation of bordeaux mixture, chemical hydrated lime for, 571–572
Presprinkling, in potato seed treatment, 793–814
Prevention of seedling diseases of sugar beets, 259–296
PRIODE, C. N., see FROMME, F. D.
Prunus spp. hosts of Exoascus cerasi, 22
Pseudomonas, pisi, causal organism of bacterial blight of pea, 611–627; pisi, host range of, 623–625; pisi, isolation of, from pea, 613–614; pisi, overwintering and seed transmission of, 614–619; tumefaciens, strains of, 46–47
Puccinia, coronata, cytological study of, on Banner and Cowra oats, 748; coronata avenae, physiologic specialization in, 783–790; dispersa, 821; emiliae, overwintering, 494–495; glumarum, 821; graminis, 822; graminis, nature of resistance of Berberis spp. to, 95–114; graminis tritici, 815; graminis tritici, chemical investigations of colored compounds in normal rust and in mutants, 720–723; graminis tritici, color mutations of, 711–725; graminis tritici, inheritance in wheat of reaction to physiologic forms of, 573–609; helianthi, effect of chemicals on germination, 496; sorghi L., course of infection in resistant and susceptible lines, 816–819; sorghi Schw., the nature of resistance in Zea mays L. to, 815–825; sorghi L., physiologic forms, 815–825; triticina, 822; triticina, effect on seed production, 40; xanthii, overwintering, 494–495
PURDY, HELEN A., Multiplication of the virus of tobacco mosaic in detached leaves, 58
Purification of the virus of tomato mosaic, 744
Pythium, a peculiar type of, 55
Pyrenomycete, causing coffee, root disease, description of, 150
Pythiacystis citrophthora, enzymes of, 747
Pythium, aphanidermatum, 43; debaryanum, cause of damping-off and root-rot, compared with P. ultimum, 54–55; spp. cause of seedling disease of sugar beet, 263; ultimum, on cabbage heads, comparison with P. debaryanum, 54–55

Quarantine, relation to forest pathology, 13

Radish, black-root, caused by **Aphanomyces raphani** n. sp., 43
RAEDER, J. M., and C. W. HUNGERFORD, Dust treatments for the control of oat smut in Idaho, 569–570; Seed treatment control of Rhizoctonia of potatoes in Idaho, 793–814
RAMSEY, G. B., and ALICE A. BAILEY, The development of nailhead spot of tomato during transit, 43–44
RANKIN, W. H., Mosaic of red and black cultivated raspberries, 46
Raspberry, Gloeosporium blight of, 769–774; mosaic, 46
RATHBUN-GRAVATT, ANNIE, A witches' broom of introduced Japanese cherry trees, 19–24
RATLIFFE, G. T., see MARTIN, J. H.
RAWLINS, T. E., Celery experiments, 746
Reaction, to stem rust in crosses between Marquis, Kota, and Kanred wheats, 595–601
Red cedar, western, leaf blight of, 11
Rehmiellopsis bohemica, cause of needle and twig blight of firs, 9
Relation of insects and weather to the development of heart rot of celery, 663–667
Report, of eighteenth annual meeting of American Phytopathological Society, 343–352; of eleventh annual meeting of Pacific Division of American Phytopathlogical Society, 745
Resistance, nature of, in Zea mays L. to Puccinia sorghi Schw., 815–825; of corn to scutellum rot, factors in, 460–469; of corn to smut, 42; relation between morphological characteristics of Berberis spp. and infection by Puccinia graminis, 98–113
Resistant variety, of cucumbers to mosaic, 48

Review of "Citrus diseases and their control," 203–204; of Japanese textbook on plant pathology, 37; of "Manual of plant diseases," 341–342
Rhabdocline pseudotsugae, cause of needle cast of Douglas fir, 7
Rhizoctonia, of potatoes, seed treatment control, in Idaho, 793–814; solani, control in coniferous seed-beds, 51–52; sp., damping-off due to, 501; spp. cause of seedling disease of sugar beets, 263; spp., causing scutellum rot of corn, 453–459
RHOADS, ARTHUR S., Clitocybe root-rot of trees and other woody plants in Florida, 56–57; Investigations on citrus "blight," wilt or leaf-curl in Florida, 58–59
Rind disease of sugar cane, 152
Ring spot of sugar cane caused by Leptosphaeria sacchari van Breda de Haan., 153; of tobacco, an infectious disease of unknown cause, 321–328
ROBERTS, JOHN W., "Target canker" of apples and pears, 735–738
RODENHISER, H. A., and E. C. STAKMAN, Physiologic specialization in Tilletia levis and Tilletia tritici, 247–253
ROLFS, P. H., Review of Fawcett and Lee's book "Citrus diseases and their control," 203–204
Root disease, of coffee, 150; of sugar cane, non-pathogenic causes, 70–82; relation of soil conditions to, 70–82; knot, on Hibiscus in Java, 26; rot, Clitocybe of trees and other woody plants in Florida, 56–57; rot of cotton, caused by Fusarium moniliforme, 227–238; rot of pea, 133–134; rot, Pythium debaryanum the cause of, 54–55; rots, of conifers, 11
ROSE, D. H., and L. F. BUTLER, Pleospora rot of lemons and apples, 47; Relation of storage temperature to lag in growth of fungous cultures, 55
Rose mildew, caused by Sphaerotheca pannosa (Wal.) Lev., 201
Roselle, diseases of in Java, 25
Rosellinia bunodes, fungus causing root disease of coffee, 151
Rot, butt, of Diospyros virginiana, 339–340; heart, of celery, 663–667; Hypochnus of apples, increasing prevalence of, 743–744; internal of caprified figs, 199; on cotton roots caused by Fusarium moniliforme, Sheld., 227–238; on scutellum of corn, 449–471; Pleospora, of lemons and apples, 47; storage of cranberries, 649–655
Rust, blister, of white pine, 10; cedar, varietal susceptibility of apple varieties to, 541–543; crown, cytological study of crown rust, 748; fixing and staining of corn, 816; germination tests with teliospores of, 491–498; leaf, effect on seed production of wheat, 40; reaction, in crosses between Marquis, Kota, and Kanred wheats, 595–601; stem, color mutations in, 711–725
RUTTLE, MABEL L., and W. P. FRASER, A cytological study of Puccinia coronata Cda. on Banner and Cowra 35 oats, 748

Saccharum officinarum, zonate foot rot of, 83–94
Sand burn, of pecan seedlings, 657–661
Saprophyte, use of term, 409–411
Scab, apple, infection studies, 45; citrus, caused by Sphaceloma fawcetti, 153; common, of potato, 297–314; of apple, control, 45; of wheat, susceptibility of wheat varieties and hybrids to, 40–41
Scaptomyza graminum, agent of inoculation of celery heart rot, 663–667
SCHERER, C. M., Tree injection for control of fungous diseases and insect pests, 51
SCHMIDT, ROBERT, see POOLE, R. F.
SCHNEIDERHAN, F. J., Biography of FRANK J. PIEMEISEL, 135–136; Black walnut (Juglans nigra L.) as a cause of the death of apple trees, 529–540
Sclerospora graminicola (Sacc.) Schroet., a study of, 57
Sclerotinia americana, Fusarium and Trichoderma on, 791–792; fructicola, fungi on, 791–792; two fungi on apothecia, 791–792
Sclerotium, delphinii, on larkspur, 240–242; delphinii, pathogenicity of, 242–244; disease of larkspur, 239–245;

rolfsii, growth on media, 418–419; rolfsii, metabolic products, 428–434; rolfsii, on Hibiscus in Java, 25–27; rolfsii, physiology and parasitism of, 53, 417–448; rolfsii, relation of H-ion concentration to growth, 420–428; rolfsii, relation to host tissue, 442–443; rolfsii, temperature relations, 419–420; rolfsii, toxicity of culture solutions, 434–441

Scutellum rot disease of corn, 449–471

Seed-beds, coniferous, control of damping-off in, 51–52

Seed, barley, dehulling with sulphuric acid to induce infection with covered smut, 747–748; certification, plant disease control by, 40; coat of corn, permeability to mercury compounds, 51; lots, differences in liability to pale dwarf of peanuts, 220–221; production, of wheat, effect of leaf rust on, 40; treatment, control of Rhizoctonia of potatoes in Idaho, 793–814; treatment, effect on growth, yield, and disease control in vegetable crops, 44; vigor, relation to pale dwarf of peanuts, 222

Seed-piece transmission of potato blackleg, 155–160

Seedling diseases of sugar beets, prevention of, 259–296; sand burn of pecan, 657–661

Semesan, dust, for control of Rhizoctonia of potatoes, 794–814; dusts, for control of barley stripe, 176–178

Separation of fern leaf from mottling in tomato mosaic, 57–58

Septoria lycopersici Speg. infecting tomato leaves, 153

Serological differentiation of Bacterium campestre from Bact. phaseoli, Bact. phaseoli sojense, and Bact. flaccumfaciens, 53–54; studies of Bacterium phaseoli, Bact. phaseoli sojense, and Bact. flaccumfaciens, 54

Setaria viridis, downy mildew on, 57

Sexual fusions, observations on, in Ustilago zeae, 831–832

S. F. A., dusts, for control of barley stripe, 176–178

SHAPOVALOV, MICHAEL, Inoculation experiments with western yellow tomato blight in relation to environmental conditions, 746

SHARP, C. G., Correlation of virulence and acid agglutination of a smooth and a rough strain of Bacterium phaseoli sojense, 49; Serological and physiological studies of Bacterium phaseoli, Bact. phaseoli sojense, and Bact. flaccumfaciens, 54; see LINK, GEO. K. K.

SHERBAKOFF, C. D., A modified method of delinting cotton seed with sulfuric acid, 189–193; Photographing living conidia mounted on agar, 195–198; Powdery mildew of muskmelon, 414–415

S. I., dust, for control of barley stripe, 176–178

Silicofluoride of sodium, theory to account for its bactericidal action and the lack of injury to host tissues, 50

Silver fir aphid, 9

SKORIC, VLADIMIR, Bacterial blight of pea: overwintering, dissemination, and pathological histology, 611–627

SMITH, ERWIN F., biography of, 675–688

Smut, covered, dehulling barley seed with sulphuric acid to induce infection with, 747–748; kernel, of sorghum, control, 52; loose kernel, on Feterita, 338–339; loose of barley, strains and host relationships, 42; millet, control, 52, 739–741; of oats, dust treatment for control of, in Idaho, 569–570; of oats, effects of soil moisture, soil temperature, and dehulling on, 31–36; resistance in corn, 42; stinking, of wheat, distribution of, 167–174

Sodium silicofluoride, theory to account for its bactericidal action and the lack of injury to host tissues, 50

Soil, chemical treatments of, for control of damping-off fungi, 499–505; conditions of, in relation to root disease of sugar cane in Cuba, 70–82; moisture, effect on smut infection of oats, 31–36; moisture, relation to citrus blight, wilt, or leaf-curl, 58–59; relation between cotton wilt and, 759; temperature, effect on smut infection of oats, 31–36; treatments for the control of damping-off in coniferous seedbeds, 51–52; types in relation to zonate

foot rot of sugar cane, 86; muck, correction of with copper, 49–50
Sooty mold of coffee, caused by Capnodium brasilense Pat. and Maubl., 152
Sorghum kernel smut, control with sulphur and copper carbonate dusts, 52
SOROKIN, HELEN, Phenomena associated with the destruction of the chloroplasts in tomato mosaic, 363–379
SPAULDING, PERLEY, A serious disease of birches, 59
Sphaceloma fawcetti, Jenkins, causal organism of citrus scab, 153
Sphacelotheca, cruenta, on Feterita, 338–339; fulginea, on muskmelon, 414–415; pannosa (Wal.) Lev. cause of rose mildew, 201
Spore germination, color mutants, Puccinia graminis tritici, 713–715
Spores, toxicity of dust fungicides to, 315–319
SPRAGUE, RODERICK, see LINFORD, MAURICE B.
Spraying, effect of early, on cucumber wilt and mosaic diseases, 473–481; effect on keeping quality of citrus fruits, 47; to prevent apple blotch cankers, 185
Sprays, oil, fungicidal value of, 201–202
Spruce, Norway, root rot of, 11
Spur blight, of raspberry, differentiated from Gloeosporium blight, 769–770
Squash, curley-top of, 48–49
Staining, of corn rust material, 816
STAKMAN, E. C., and J. J. CHRISTENSEN, Heterothallism in Ustilago zeae, 827–834; see CHRISTENSEN, J. J.; see RODENHISER, H. A.
Statistical studies, spores, color mutants, Puccinia graminis tritici, 715–719
Stem-end rot, of citrus fruits, 47
STEVENS, F. L., and P. A. YOUNG, Use of terms saprophyte and parasite, 409–411
STEVENS, NEIL E., and HENRY F. BAIN, Storage rots of cranberries in the 1926 crop, 649–655
STEWART, DEWEY, see COONS, G. H.
Stilbum leaf spot, outbreak of, in Cundinamarca, 147
Stinking smut, of wheat, distribution of, 167–174
Stripe disease of barley, dusts for control of, 157–179
Striped cucumber beetle, control, 473–481
Storage rots, of cranberries, 649–655; temperature, relation to lag in growth of fungous cultures, 55
Strains, see physiologic specialization
Strawberry, preliminary studies on witches' broom of, 329–335
Studies on the control of millet smut, 739–741; on the scutellum rot disease of corn, 449–471; of growth habit and rust reaction in crosses between Marquis, Kota, and Kanred wheats, 573–609
Study of fungous flora of nodal tissues of corn plant, 563–568
SUEMATU, N., Text-book on plant pathology, review of, 37
Sugar beet, curly top of, in Argentine, 407–408; curly-top virus, the cause of western yellow tomato blight, 39; ecological studies of curly top of, 747; prevention of seedling diseases of, 259–296
Sugar cane, effects of mosaic on content of cell of, 57; eye spot of, 315; "gomosis," 148–150; mosaic, corn mosaic of Hawaii distinct from, 41; rind disease, caused by Melanconium sacchari Massee, 152; ring spot disease of, 153; root disease of, in Cuba, 61–82; zonate foot rot of, 83–94
Sulphur, as a control agent for common scab of potatoes, 297–314; dusts for control of sorghum kernel smut and millet smut, 52
Sulphuric acid, dehulling barley seed with, to induce infection with covered smut, 747–748; used for delinting cotton seed, 189–193
Summer meeting, American Phytopathological Society, 415, 836
Susceptibility of different hosts to Aphanomyces euteiches, 133–134; apple varieties to cedar rust, 541–543; of wheat varieties and hybrids to wheat scab in Minnesota, 40–41
Sweet pea fasciation, a form of crown-gall, 29–30; potatoes, Fusarium wilt of, 42–43

Symptoms, of dwarf disease of blackberries, 633–639; of sand burn of pecan seedlings, 657–658

TAKAHASHI, TAKAMICHI, A Sclerotium disease of larkspur, 239–245

Taphrina aurea, strains on Populus spp., 12

"Target canker" of apple, relation to tree vigor, 738

Teliospores of rust, germination tests, 491–498

Temperature, leaf, in relation to tip burn of lettuce, 44–45; relations, inoculations, tobacco mosaic on potatoes, 731–732; relations, Sclerotium rolfsii, 419–420; relations to growth of Penicillium spp. and citrus fruit decay, 746–747; soil, effect on smut infection of oats, 31–36; storage, relation to lag in growth of fungous cultures, 55

Tenebriodes mauritanicus (L.) a leaf-eating insect, 145

THOMAS, H. E., Some chemical treatments of soil for the control of damping-off fungi, 499–505

Thread blight disease caused by Corticium koleroga on citrus and pomaceous plants, 689–709

Tillantin "C," for control of barley stripe, 175

Tilletia, distribution of species of, 167–174; favorable temperature range for infection by species of, 168–169; laevis, a study of distribution of T. tritici and, 167–174; laevis, occurrence of, 168–173; laevis, physiologic specialization in, 247–253; tritici and T. laevis, a study of distribution of, 167–174; tritici, occurrence of, 168–173; tritici, physiologic specialization in, 247–253

Tip burn, of lettuce, relation of leaf temperature to, 44–45

TISDALE, W. H., and MARION A. GRIFFITHS, Strains of Ustilago nuda and certain host relationships, 42; C. E. LEIGHTY, and E. G. BOERNER, A study of the distribution of Tilletia tritici and T. laevis in 1926, 167–174

Tissues, nodal, of corn plant, study of fungous flora of, 563–568

Tobacco, bug, Nezara viridis, 181; barley, effect of a strain of mosaic on yield and quality of, 523–527; effects of mosaic on content of cell of, 57; inoculated with potato virus, 777–778; leaves infested by Nezera viridis (green bug) attacked by Blakeslea trispora, 181; mosaic, 152; mosaic disease of, 513–527; mosaic, effect of age on virus, 515–517; mosaic, effect of a strain of, on yield and quality of burley tobacco, 523–527; mosaic, from commercial and cured leaf tobaccos, 49; mosaic on potatoes, 727–734; mosaic virus, multiplication in detached leaves, 58; "natural leaf," chewing, mosaic in, 517; ringspot of, 321–328; commercial, and cured leaf as source of tobacco mosaic, 49, 513–522

Tomato, leaves infected with Septoria lycopersica, 153; mosaic, destruction of chloroplasts in, 363–379; mosaic, purification of virus, 744; mosaic, separation of fern leaf from mottling, 57–58; nailhead spot of, development during transit, 43–44; western yellow blight caused by beet curly-top virus, 39; western yellow blight, 746; yellows, 746; control of damping-off, 500–503

TORO, RAFAEL A., see CHARDON, CARLOS E.

Toxicity, culture solutions of Sclerotium rolfsii, 434–441; of dust fungicides, a method of testing in vitro, 315–319

Trametes rigida, 137

Transmission, of curly top of sugar beet by Agallia stictiocollis, 407–408; of dwarf disease, 639–646; seed, of bacterial blight of pea, 614–619; seed piece, of potato blackleg, 155–160

Tree injection for control of fungous diseases and insect pests, 51

Trees, fruit, treatment of lime-induced chlorosis in, 745–746; in Florida, Clitocybe root-rot of, 56–57

Trichoderma lignorum or koningi on Sclerotinia americana, 791–792

Trifolium, hybridum, incarnatum, medium, pratense, repens, leafhopper injury of, 399–404

TRUE, RODNEY H., Biography of ERWIN F. SMITH, 675–688

Turf, fungicidal control of brown-patch of, 50–51
Twig, and needle, blight of firs, 9; and needle, blight of pines, 9
Type culture collection, American, 672

Uredo chrysanthemi, 821
Uromyces, appendiculatus, bean rust, 152; spp., effect of potassium dihydrogen phosphate on germination of teliospores, 495–496
Uspulun, for control of damping-off of tomatoes, 499–505
Ustilago, hordei, dehulling barley seed with sulphuric acid to induce infection with, 747–748; nuda, strains and certain host relationships, 42; violacea (Pers.) Fuckel., 827; zeae, behavior of individual physiologic forms, 827–831; zeae, behavior of physiologic forms in various combinations, 827–831; zeae, heterothallism in, 827–834; zeae, observation on sexual fusions, 831–832; zeae, possible origin of physiologic forms, 833
Ustulina, maxima (Weber) v. Wettst., comparison with U. zonata and U. vulgaris Tul., 143; vulgaris Tul., 143; zonata (Lev.) Sacc., a fungus with wide host range in tropics, 145; zonata (Lev.) Sacc. associated with Ganoderma pulverulentum, 142

Vaccinium macrocarpon, storage rots of, 649–655; parvifolium, Exobasidium on, 207–211
VALLEAU, W. D., and E. M. JOHNSON, Commercial tobaccos and cured leaf as sources of tobacco mosaic, 49; Commercial tobaccos and cured leaf as a source of mosaic disease in tobacco, 513–522; The effect of a strain of tobacco mosaic on the yield and quality of burley tobacco, 523–527
VAN HALTERN, FRANK, see MELHUS, I. E.
Variation of strains of Alternaria solani isolated from lesions on potato tubers, 56
Variety, of cucumbers resistant to mosaic, 48
Venturia inaequalis (Cke.) Wint., effects of fungous extracts upon the initiation and growth of the perithecia of, 835–836
Vicia, angustifolia, dasycarpa, ervilea, fulgens, gigantea, monatha, pannonica, sativa, hosts of Aphanomyces euteiches, 133
Viroses of Solanum tuberosum, uniformity of nomenclature for, 161–165
Virulence, correlation in strains of Bacterium phaseoli sojense, 49
Virus, disease of blackberries, 646; of tobacco mosaic, multiplication of in detached leaves, 58; of tomato mosaic, purification of, 744; of potato, on peppers, 775–782; tobacco mosaic, effect of age on, 515–517

Walnut, black, as the cause of the death of apple trees, 529–540
WALTON, RICHARD CHAMBERS, In memoriam, 353
Watermelon, wilt infection studies, 47–48
WATSON, ELIZABETH, see LEWIS, I. M.
"Wa-Wa" dust, for control of barley stripe, 176–178
Weather, relation of, to the development of heart rot of celery, 663–667
WEBB, ROBERT W., Certain factors influencing the development of the mosaic disease in winter wheat, 41
Weed tree, Dichrostachys nutans, in Cuba, with remarks on its pathology, 137–146
WEIMER, J. L., A wilt disease of alfalfa caused by Fusarium sp., 337–338
WEIR, JAMES R., Butt rot of Diospyros virginiana caused by Polyporus spraguei, 339–340; the problem of Dichrostachys nutans, a weed tree, in Cuba, with remarks on its pathology, 137–146
WELCH, CATHERINE GRAHAM, see DUFF, G. H.
WELLENSIEK, S. J., The nature of resistance in Zea mays L. to Puccinia sorghi Schw., 815–825
Western yellow tomato blight, caused by beet curly-top virus, 39; yellow tomato blight in relation to environmental conditions, 746
WESTON, W. H., see COLLEY, REGINALD, H.
Wheat, effect of leaf rust on seed production of, 40; inheritance of growth habit

and rust reaction, 573–609; mosaic disease of, 41; scab, susceptibility of wheat varieties and hybrids to, 40–41
White pine blister rust, 10
WIANT, J. STEWART, Soil treatments for the control of damping-off in coniferous seed-beds, 51–52
WILSON, ALBERT, Preliminary report on the gardenia bud drop, 671–672
WILSON, E. E., Effects of fungous extracts upon the initiation and growth of the perithecia of Venturia inaequalis (Cke.) Wint. in pure culture, 835–836; see KEITT, G. W., 45
Wilt, blue-stem, of raspberry, differentiated from Gloeosporium blight, 769–770; citrus, in Florida, 58–59; cucumber, control, 473–479; disease of alfalfa caused by Fusarium sp., 337–338; Fusarium, of sweet potatoes, 42–43; of cotton and its control, 749–767; watermelon, infection studies, 47–48
WINGARD, S. A., see FROMME, F. D.
Witches' broom, of introduced Japanese cherry trees, 19–24; of strawberry, preliminary studies on, 329–335
WOLF, FREDERICK A., and WALTER J. BACH, The thread blight disease caused by Corticum koleroga (Cooke) Höhn., on citrus and pomaceous plants, 689–709
WOODROOF, NAOMI CHAPMAN, A disease of cotton roots proluced by Fusarium moniliforme Sheld., 227–238
Woody plants, in Florida, Clitocybe root-rot of, 56–57

X-bodies in the cells of "mosaic diseased" and "dwarfed" dahlias, 52
Yellow dwarf disease of potatoes, on peppers, 775–782
Yellowing, "amarillamiento," root disease of coffee, 150; of alfalfa, caused by leafhoppers, 39
Yellows, of clover, due to leafhoppers, 58; of tomato, 746
YOUNG, H. V., Summer meeting of American Phytopathological Society, 415
YOUNG, P. A., see STEVENS, F. L.
YOUNG, V. H., Varietal susceptibility of Ada Red and certain other apple varieties to cedar rust, with special reference to twig infections, 541–543

Zea mays, see corn
ZELLER, S. M., Dwarf of blackberries, 629–648; Preliminary studies on witches' broom of strawberry, 329–335
Zonate, foot rot of sugar cane, 83–94